

Review

Ellipsoid zone on optical coherence tomography: a review

Lingwei William Tao MBBS,¹ Zhichao Wu PhD,¹ Robyn H Guymer PhD FRANZCO^{1,2} and Chi D Luu PhD^{1,2}

¹Centre for Eye Research Australia, Royal Victorian Eye and Ear Hospital, and ²Department of Surgery (Ophthalmology), The University of Melbourne, Melbourne, Victoria, Australia

ABSTRACT

Emergence of the high-resolution optical coherence tomography has allowed better delineation of retinal layers, and many of the anatomical correlations of these layers have now been agreed upon. However, some anatomical correlates still remain contentious, such as the second hyper-reflective band, which is now termed ellipsoid zone. Despite the lack of consensus of the actual origin of the ellipsoid zone, there has been much interest in evaluating its integrity and intensity in different disease processes. This review paper aims to provide an overview of the ellipsoid zone and its clinical and research applications.

Key words: diagnostic study, diagnostic technique, retinal disease, retinal imaging.

INTRODUCTION

Since its first emergence two decades ago, optical coherence tomography (OCT) has revolutionized our ability to visualize the retina *in vivo* and to identify the anatomical correlates of pathological changes in ocular diseases. As such, it has facilitated in the clinical management of various retinal and optic nerve conditions.¹

Optical coherence tomography has demonstrated its potential for identifying novel markers of different disease states as it allows detection of ultrastructural

changes in the retina when there is no clinical correlate to visualize. Improved resolution with advances in OCT technology has allowed the delineation of individual retinal layers providing more information to increasing our knowledge of specific subclinical changes in many retinal diseases such as drusen ultrastructure in age-related macular degeneration (AMD),² optic nerve head and retinal nerve fibre layer in glaucoma.^{3–5} The emergence of the spectral domain OCT (SD-OCT) has allowed better delineation of retinal layers such as the vitreoretinal interface, inner and outer retina, retinal pigment epithelium (RPE) and choroid. Many of these layers have been universally agreed upon; however, some anatomical correlates remain contentious,⁶ such as the second hyper-reflective band. This band was previously believed to represent the junction between the inner and outer segments (IS/OS junction) of the photoreceptors.^{1,7,8} More recent evidence suggests that the second hyper-reflective band corresponds with the anatomical location of the ellipsoid portion of the photoreceptors inner segment.^{1,7,8} In an effort to come to a consensus on the anatomical correlates with OCT description of each layer, an international panel with expertise in vitreoretinal diseases and imaging met in 2014 to access available evidence and provide recommended OCT terminology.⁶ The international nomenclature consensus panel recommended the second hyper-reflective band be termed as the ellipsoid zone (EZ), as it avoids attribution of the OCT feature specifically to a single anatomic structure until more definitive evidence becomes available, and this

■ **Correspondence:** A/Professor Chi D Luu, Macular Research, Centre for Eye Research Australia, Level 8, 32 Gisborne Street, Melbourne, Vic. 3002, Australia. Email: cluu@unimelb.edu.au

Received 21 August 2015; accepted 3 November 2015.

Competing/conflicts of interest: None.

Funding sources: This research was supported by the National Health and Medical Research Council (NH&MRC) project grant (1084081) and BrightFocus Foundation. Centre for Eye Research Australia (CERA) receives Operational Infrastructure Support from the Victorian Government and is supported by a NHMRC Centre for Clinical Research Excellence Award (#529923).

paper will use this terminology throughout. Despite the lack of consensus of the actual origin of the EZ, there has been much interest in its appearance particularly evaluating its integrity and intensity in different disease processes.^{9–24} In this review paper, we provided an overview of the EZ and its clinical and research applications.

Optical coherence tomography

Optical coherence tomography is a non-invasive imaging modality that produces high-resolution cross-section images of the retinal microanatomy.²⁵ The principle of OCT is based on low-coherence interferometry, which analyses the difference in back-scattered light from tissues.^{25,26} OCT utilizes a broadband light source, which is first split into a reference and sample beam. With the back-scattered beam from the retina generating an interference pattern with the reference beam, a reflectivity *versus* depth profile can be constructed.^{25,27} Initially, OCT was only able to provide axial resolution of around 10 to 20 μm , which did not allow for the further differentiation of the neurosensory retina and the RPE, which appeared as one single highly reflective band.²⁸ Earlier generation time-domain OCT (TD-OCT) like Stratus OCT (Carl Zeiss Meditec, Inc., Dublin, CA, USA) employed time-domain method that allowed a 400 A-scans per second scan rate with an axial resolution of 8 to 10 μm .²⁷ The newer generation SD-OCT, such as the Spectralis (Heidelberg Engineering GmbH, Heidelberg, Germany) and Cirrus OCT (Carl Zeiss Meditec, Dublin, CA, USA), employs an interferometer and spectrometer, which analyses back-scattered light interference pattern simultaneously through Fourier transform algorithm.^{29,30} This innovative technique enabled an axial resolution of 5 to 7 μm , faster scan rates at 20 000 to 52 000 A-scans per second and better signal-to-noise ratio compared with TD-OCT.^{26,29–32} The improved delineation of different retinal layers makes this modality ideal for clinical diagnosis and management of ocular diseases (Fig. 1).

Retinal ultrastructures visualized by OCT

Based upon early TD-OCT, structural correlations with the OCT images were first reported. In 1991, a TD-OCT was capable of producing a single A-scan in 1.25 s with an axial resolution of 17 μm .²⁵ The inner retina hyper-reflective zone was reported to represent the nerve fibre layer, but the outer retina hypo-reflective zone was not further described. Using different animal models, comparisons between OCT images and histological data were considered. Using Macaca mulatta, Toth *et al.*³³ found that the outer

nuclear layer and the inner and outer segments of the photoreceptors had low relative reflectivity. In a study using chickens, Huang *et al.*³⁴ showed a hyper-reflective zone in outer retina, which they speculated to be associated with the increase in local refractive index at the photoreceptor inner segment because of the presence of mitochondria.

In early 2000, the ability to perform higher resolution OCT allowed various layers of the outer retina to be delineated. Three to four distinct hyper-reflective bands on SD-OCT can be identified in the human outer retina.³⁵ The first band is the innermost and the least intense of the four hyper-reflective bands, which is thought to originate from the external limiting membrane (ELM).³⁶ The second band is now referred to as the EZ.⁶ The histological correlation of the third band is still debated as it has been attributed to the cone outer segment tips,³⁷ Verhoeff's membrane^{35,38} or as the intermediate line but is referred to as the interdigitation zone.³⁹ The fourth band is thought to represent the RPE with contribution from both Bruch's membrane and the choriocapillaris.⁴⁰

Origin and significance of the ellipsoid zone

The second hyper-reflective band has been traditionally ascribed to the inner/outer segment junction of the photoreceptors.⁴¹ However, recent studies suggest that this band is anatomically correlated with the photoreceptor inner segment ellipsoid (ISE)^{1,7} and is referred to in the international nomenclature consensus as the EZ.⁶ The photoreceptor ISE has a high refractive index, leading to increased back-scattering of light that translates to the bright appearance of the EZ.^{42–45} The photoreceptor ISE is densely packed with mitochondria and is thus important for the photoreceptor health and function. Hence, the ability to link a signal from this band with a diseased state would be interesting and suggestive of mechanisms at play in disease causation.

Using adaptive optics, Jonnal *et al.*⁴⁶ argued that the EZ is both too thick and proximally located to be generated by the ellipsoids of the inner segments. However, only the foveal cone was imaged in that study, which may explain the inconsistent findings with the earlier reports. Although the exact anatomical correlation of the EZ remains disputed, its roles in the clinic and in research are still relevant because change in its integrity and intensity has been correlated with different retinal disease processes.^{9,21,40,47–49}

Changes in the ellipsoid zone in different disease processes

Changes in the EZ are determined based upon its integrity or intensity. The integrity of the EZ can be characterized broadly into three categories of presence,




Figure 1. An example of a spectral domain optical coherence tomography of a normal eye.

disruption or absence. More recently, a quantitative approach of determining the relative intensity of the EZ has been advocated. The relative intensity of the OCT EZ can be expressed as a ratio of the EZ to the ELM.^{10–12} The intensity of the ELM layer is used as the reference value to control for the variation in the brightness of different OCT scans because the ELM is a non-neural layer, does not alter its intensity with age or the stage of degeneration and as it has a consistent intensity across the retina.¹⁰ The inner nuclear layer (INL) has also been investigated for the use as a reference layer because it has a larger area compared with the ELM, which could have made the manual measurements of the intensity easier. At present, the ability to determine the intensity of the EZ nor the ratio with the ELM is not automated so that time-consuming manual examination is required.^{10–12,17} The EZ integrity and intensity have been studied in various retinal conditions.

Retinal and macular degenerative conditions

Changes in EZ intensity have been reported in retinal degenerative conditions. Hood *et al.*¹⁷ investigated the intensity of the EZ in cone dystrophy, a condition known to have reduced number of functioning cones, and found that the disease produced a less intense signal from the EZ compared with healthy controls. Sundaram *et al.*¹² investigated the EZ intensity in patients with achromatopsia and its implication in establishing measures to determine patients suitable for gene therapy. They found that those without any disruption in the EZ, the mean intensity ratio of the EZ was significantly lower in achromatopsia compared with controls, but there was no correlation between intensity and retinal sensitivity measured

by microperimetry.¹² Furthermore, a case report of peripheral cone dystrophy has found that the reduction in the OCT EZ intensity is to be correlated with the reduction in cone photoreceptors in the corresponding area imaged by adaptive optics.⁵⁰

In our own studies, we found that patients with the early stages of AMD exhibit reduced relative intensity of the EZ when compared with healthy controls across different retinal eccentricities, except at the fovea.¹¹ We have also observed a correlation between the reduction in relative intensity of the EZ with delayed multifocal electroretinogram (mfERG) implicit time, but not with the mfERG amplitude.¹⁰ In a separate study in eyes with intermediate AMD, we observed the recovery of the EZ integrity following drusen regression without evidence of nascent geographic atrophy (nGA). The recovery of the EZ integrity was associated with the improvement in retinal function as measured by the mfERG (Fig. 2).

Interestingly, the age-related maculopathy susceptibility 2 (*ARMS2*) gene, implicated in AMD, is considered by some to have a role in mitochondrial homeostasis, impacting upon the cellular apoptosis pathway of the cones, although there is not universal agreement on the role of this gene.⁵¹ A decrease in the EZ intensity may be reflecting the reduction in healthy or functional mitochondria in the photoreceptor inner segment, which could translate to abnormal photoreceptor function. Therefore, a quick retinal scan with OCT could potentially detect AMD patients with *ARMS2* genotypes and provide a surrogate measure of retinal function.

Disruption of EZ has also been reported in AMD eyes with reticular pseudodrusen (RPD)^{14,20,52} and it has been shown to be associated with the reduction of cone density.¹⁴ In a longitudinal study in eyes with RPD, Spaide *et al.* found that eyes progressed




Figure 2. Optical coherence tomography (OCT) images and multifocal electroretinogram (mfERG) responses from an intermediate age-related macular degeneration eye. OCT scans show drusen regression and change in the ellipsoid zone (EZ) integrity at the 12-month visit and the recovery of the EZ integrity at the 24-month visit. The recovery of the EZ integrity is associated with the improvement in the mfERG implicit time, particularly at the central and temporal retina.

to atrophy were associated with loss of photoreceptor length, disruption of EZ band or decrease EZ intensity, and visual loss.⁵³ In eyes with geographic atrophy (GA) secondary to AMD, it has been demonstrated that retinal areas with disrupted EZ have a greater risk for progressing to dense scotoma compared with areas with intact EZ.⁵⁴ Several retrospective studies have investigated the EZ changes in patients treated for neovascular AMD,^{55,56} with the presence of the EZ associated with better final visual acuity following photodynamic therapy than in those where the zone was missing.⁵⁷

Inflammatory diseases

Disruption of the EZ has been observed (Fig. 3) in patients with multiple evanescent white dot syndrome (MEWDS).^{9,58,59} Hashimoto *et al.*⁵⁹ described diffuse disruption of the EZ in the acute stage, with restoration of the EZ in the intermediate stage and its complete restoration in the late stage of MEWDS. The disappearance of the EZ is initially thought to be related to the structural change in the photoreceptor secondary to disease process, which alters its refractive property and makes it invisible on OCT. However, it is noted that the band reappears as the disease evolves suggesting perhaps that the disruption is likely a reflection of altered refractive characteristics,

which resolve as the inflammation resolves, rather than a permanent loss of cells or their function.⁹

Other inflammatory conditions associated with changes in EZ integrity include acute posterior multifocal placoid pigment epitheliopathy (APMPPE), punctate inner choroidopathy (PIC) and acute zonal occult outer retinopathy (AZOOR). In APMPPE, the EZ is often disrupted in the acute stage of the disease,⁶⁰ but EZ disruption can also be seen after healing in areas that initially appeared normal on OCT.⁶¹ EZ disruption is also observed in PIC,^{62,63} and it is usually associated with the focal elevation of the RPE.⁶⁴ In AZOOR, the EZ is often irregular or disrupted at the lesion sites or retinal areas with abnormal mfERG.^{65,66} Recovery of the EZ integrity has also been reported in some AZOOR patients when retinal function and enlarged blind spot improved.⁶⁷

Other retinal conditions

The integrity of the EZ has been found to be an important indicator of visual outcomes,⁶⁸ and its normal appearance is associated with better post-operative visual acuity in idiopathic macular hole,^{22,69} in epiretinal membrane^{70–72} and in retinal detachment.^{73–75} Other retinal diseases, where the integrity of the EZ was found to correlate with visual outcome, include macular telangiectasia,⁷⁶ retinal vein occlusion^{77,78} and central serous chorioretinopathy.^{79,80}




Figure 3. Fundus autofluorescence and optical coherence tomography (OCT) images of an eye with multiple evanescent white dot syndrome. The OCT image shows the disruption of the ellipsoid zone (arrows) at the lesion.

The association between the EZ integrity and visual outcome has also been demonstrated in diabetic macular oedema. The extent of the EZ disruption was found to be a predictor of visual outcome in diabetic macular oedema.^{81,82} It has been shown in a retrospective interventional study that the shorter length of disruption in the EZ correlated with better visual outcome in patients with diabetic macular oedema treated with intravitreal triamcinolone.⁸³ In a study of diabetic retinopathy where there is angiographic evidence of macular non-perfusion, the authors have reported corresponding SD-OCT demonstrating both disruption of integrity and reduced intensity of the EZ.⁸⁴

Changes in EZ integrity have also been demonstrated in ocular infectious conditions such as syphilis⁸⁵ and dengue fever-associated maculopathy.⁸⁶

En-face OCT imaging techniques

En-face OCT imaging was first reported in literature in 1997.⁸⁷ Wanek *et al.*⁸⁸ have identified the EZ en-face images and demonstrated non-uniform variation in texture, which the authors attributed to limited fixation stability of the test subject. Rosenfeld *et al.* reported a novel use of en-face image processing algorithm for identifying areas at risk of GA progression, in their study, darker areas on the en-face slab images of the EZ correlated with areas that developed GA, suggesting that EZ disruption may represent an early defect preceding GA.⁸⁹ Although to date, there is no reported study investigating the change in the EZ intensity on en-face OCT, a volumetric analysis of the EZ or a measure of overall intensity, might have the potential to become a surrogate measure of disease severity in AMD and other retinal conditions.

Current limitations and future direction

The advancement in OCT technology has resulted in improved scan resolution, speed and signal-to-noise

ratio, which has translated into better delineation of subtle changes of retinal ultrastructures secondary to disease pathology. As this field is ever evolving because of new innovation in imaging techniques, clinicians who utilize OCT in their daily practice for clinical decision-makings must also be aware of the current limitation of this relatively new technology.

Although SD-OCT is capable of resolving the different layers of the retina, it lacks the lateral resolution to resolve cellular details of each layer.⁴⁶ Newly emerged imaging techniques such as adaptive optics have been shown to combine both cellular lateral resolution and depth resolution offered by the conventional SD-OCT.^{35,41,90} Consequently, previously proposed anatomical correlates are being questioned in newer studies, and as such, we should remain open to the need to re-evaluate our current assumptions on anatomical correlates to the bands and the zones seen on OCT.

Commercially available OCTs do not display the raw linear data because of the large amount of back-scattered light captured in generating the native images. Final images are usually produced from transformation of the linear OCT data through combination of logarithmic or root transformation.⁹¹ Furthermore, the SD-OCT final output image is both dependent on and limited by the instrument's ability to resolve physical structural details, also known as the point spread function of the instrument. The axial dimension of an imaged object is therefore dictated by the axial resolution of the instrument itself. For example, an infinitely thin hypothetical reflector when imaged by an OCT would have appeared as thick as the point spread function of the instrument,⁹¹ and in case of the Spectralis SD-OCT, it would appear as a hyper-reflective band measuring at 7 μm . Therefore, the final OCT images may misrepresent actual difference in reflectivity.^{1,12}

There are always further improvements in imaging, such that now with swept source OCT, we can

visualize deep structures in the posterior pole such as the choroid and therefore allow the ability to assess differences in choroidal structure with disease.^{24,92–94} More recently, there are devices that combine OCT imaging with angiography allowing blood vessels and flow to be determined without the need for formal fluorescein and indocyanine green angiography. With ever evolving techniques, we are sure to learn more about the ultrastructure and vasculature of the retina and choroid, and their changes in different eye conditions.

CONCLUSION

The second hyper-reflective band on SD-OCT is currently referred as the EZ. Although there is still a lack of a consensus on the exact anatomical correlate of the EZ, the integrity and intensity of the EZ has been shown to be associated with the visual function and is an important indicator of visual outcomes in many retinal conditions. With a relatively quick and highly reproducible OCT image acquisition, the integrity and intensity of the EZ has the potential for assessing the health of the outer retina in clinical and research settings.

Method of literature search

References for this review were identified through a comprehensive literature search of the following electronic databases: MEDLINE, PubMed and Science Direct. In addition, articles, textbooks and thesis thought to be relevant were also selected from review of the bibliographies of those articles generated from the aforementioned search. The following keywords and combinations of these words were used in compiling the search: ‘optical coherence tomography’, ‘second hyper-reflective band’, ‘IS/OS junction’, ‘inner segment ellipsoid’, ‘integrity’ and ‘intensity’.

REFERENCES

1. Spaide RF, Curcio CA. Anatomical correlates to the bands seen in the outer retina by optical coherence tomography: literature review and model. *Retina* 2011; **31**: 1609–19.
2. Khanifar AA, Koreishi AF, Izatt JA, Toth CA. Drusen ultrastructure imaging with spectral domain optical coherence tomography in age-related macular degeneration. *Ophthalmology* 2008; **115**: 1883–90.
3. Alasil T, Wang K, Keane PA, *et al.*. Analysis of normal retinal nerve fiber layer thickness by age, sex, and race using spectral domain optical coherence tomography. *J Glaucoma* 2013; **22**: 532–41.
4. Chen TC. Spectral domain optical coherence tomography in glaucoma: qualitative and quantitative analysis of the optic nerve head and retinal nerve fiber layer (an AOS thesis). *Trans Am Ophthalmol Soc* 2009; **107**: 254–81.
5. Chen TC, Zeng A, Sun W, Mujat M, de Boer JF. Spectral domain optical coherence tomography and glaucoma. *Int Ophthalmol Clin* 2008; **48**: 29–45.
6. Staurengi G, Sadda S, Chakravarthy U, Spaide RF. International Nomenclature for Optical Coherence Tomography P. Proposed lexicon for anatomic landmarks in normal posterior segment spectral-domain optical coherence tomography: the IN*OCT consensus. *Ophthalmology* 2014; **121**: 1572–8.
7. Fernandez EJ, Hermann B, Povazay B, *et al.*. Ultrahigh resolution optical coherence tomography and pancorrection for cellular imaging of the living human retina. *Opt Express* 2008; **16**: 11083–94.
8. Lu RW, Curcio CA, Zhang Y, *et al.*. Investigation of the hyper-reflective inner/outer segment band in optical coherence tomography of living frog retina. *J Biomed Opt* 2012; **17**: 060504.
9. Yang CS, Wang AG, Lin YH, Huang YM, Lee FL, Lee SM. Optical coherence tomography in resolution of photoreceptor damage in multiple evanescent white dot syndrome. *J Chin Med Assoc* 2012; **75**: 663–6.
10. Wu Z, Ayton LN, Guymer RH, Luu CD. Relationship between the second reflective band on optical coherence tomography and multifocal electroretinography in age-related macular degeneration. *Invest Ophthalmol Vis Sci* 2013; **54**: 2800–6.
11. Wu Z, Ayton LN, Guymer RH, Luu CD. Second reflective band intensity in age-related macular degeneration. *Ophthalmology* 2013; **120**: 1307–8 e1301.
12. Sundaram V, Wilde C, Aboshiha J, *et al.*. Retinal structure and function in achromatopsia: implications for gene therapy. *Ophthalmology* 2014; **121**: 234–45.
13. Oishi A, Shimozone M, Mandai M, Hata M, Nishida A, Kurimoto Y. Recovery of photoreceptor outer segments after anti-VEGF therapy for age-related macular degeneration. *Graefes Arch Clin Exp Ophthalmol* 2013; **251**: 435–40.
14. Mrejen S, Sato T, Curcio CA, Spaide RF. Assessing the cone photoreceptor mosaic in eyes with pseudodrusen and soft Drusen *in vivo* using adaptive optics imaging. *Ophthalmology* 2014; **121**: 545–51.
15. Mitamura Y, Aizawa S, Baba T, Hagiwara A, Yamamoto S. Correlation between retinal sensitivity and photoreceptor inner/outer segment junction in patients with retinitis pigmentosa. *Br J Ophthalmol* 2009; **93**: 126–7.
16. Inoue M, Watanabe Y, Arakawa A, Sato S, Kobayashi S, Kadonosono K. Spectral-domain optical coherence tomography images of inner/outer segment junctions and macular hole surgery outcomes. *Graefes Arch Clin Exp Ophthalmol* 2009; **247**: 325–30.
17. Hood DC, Zhang X, Ramachandran R, *et al.*. The inner segment/outer segment border seen on optical coherence tomography is less intense in patients with diminished cone function. *Invest Ophthalmol Vis Sci* 2011; **52**: 9703–9.
18. Gomes NL, Greenstein VC, Carlson JN, *et al.*. A comparison of fundus autofluorescence and retinal structure in patients with Stargardt disease. *Invest Ophthalmol Vis Sci* 2009; **50**: 3953–9.
19. De Bats F, Wolff B, Vasseur V, *et al.*. “En-face” spectral-domain optical coherence tomography findings in multiple evanescent white dot syndrome. *J Ophthalmol* 2014; **2014**: 928028.

20. Curcio CA, Messinger JD, Sloan KR, McGwin G, Medeiros NE, Spaide RF. Subretinal drusenoid deposits in non-neovascular age-related macular degeneration: morphology, prevalence, topography, and biogenesis model. *Retina* 2013; **33**: 265–76.
21. Chang LK, Koizumi H, Spaide RF. Disruption of the photoreceptor inner segment-outer segment junction in eyes with macular holes. *Retina* 2008; **28**: 969–75.
22. Baba T, Yamamoto S, Arai M, et al.. Correlation of visual recovery and presence of photoreceptor inner/outer segment junction in optical coherence images after successful macular hole repair. *Retina* 2008; **28**: 453–8.
23. Aizawa S, Mitamura Y, Baba T, Hagiwara A, Ogata K, Yamamoto S. Correlation between visual function and photoreceptor inner/outer segment junction in patients with retinitis pigmentosa. *Eye* 2009; **23**: 304–8.
24. Adhi M, Duker JS. Optical coherence tomography – current and future applications. *Curr Opin Ophthalmol* 2013; **24**: 213–21.
25. Huang D, Swanson EA, Lin CP, et al.. Optical coherence tomography. *Science* 1991; **254**: 1178–81.
26. Schuman JS. Spectral domain optical coherence tomography for glaucoma (an AOS thesis). *Trans Am Ophthalmol Soc* 2008; **106**: 426–58.
27. Sull AC, Vuong LN, Price LL, et al.. Comparison of spectral/Fourier domain optical coherence tomography instruments for assessment of normal macular thickness. *Retina* 2010; **30**: 235–45.
28. Hee MR, Izatt JA, Swanson EA, et al.. Optical coherence tomography of the human retina. *Arch Ophthalmol* 1995; **113**: 325–32.
29. Leitgeb R, Hitzinger C, Fercher A. Performance of Fourier domain vs. time domain optical coherence tomography. *Opt Express* 2003; **11**: 889–94.
30. de Boer JF, Cense B, Park BH, Pierce MC, Tearney GJ, Bouma BE. Improved signal-to-noise ratio in spectral-domain compared with time-domain optical coherence tomography. *Opt Lett* 2003; **28**: 2067–9.
31. Yaqoob Z, Wu J, Yang C. Spectral domain optical coherence tomography: a better OCT imaging strategy. *Biotechniques* 2005; **39**: S6–13.
32. Gloesmann M, Hermann B, Schubert C, Sattmann H, Ahnelt PK, Drexler W. Histologic correlation of pig retina radial stratification with ultrahigh-resolution optical coherence tomography. *Invest Ophthalmol Vis Sci* 2003; **44**: 1696–703.
33. Toth CA, Narayan DG, Boppart SA, et al.. A comparison of retinal morphology viewed by optical coherence tomography and by light microscopy. *Arch Ophthalmol* 1997; **115**: 1425–8.
34. Huang Y, Cideciyan AV, Papastergiou GI, et al.. Relation of optical coherence tomography to microanatomy in normal and rd chickens. *Invest Ophthalmol Vis Sci* 1998; **39**: 2405–16.
35. Zawadzki RJ, Jones SM, Olivier SS, et al.. Adaptive-optics optical coherence tomography for high-resolution and high-speed 3D retinal *in vivo* imaging. *Opt Express* 2005; **13**: 8532–46.
36. Drexler W, Sattmann H, Hermann B, et al.. Enhanced visualization of macular pathology with the use of ultrahigh-resolution optical coherence tomography. *Arch Ophthalmol* 2003; **121**: 695–706.
37. Srinivasan VJ, Monson BK, Wojtkowski M, et al.. Characterization of outer retinal morphology with high-speed, ultrahigh-resolution optical coherence tomography. *Invest Ophthalmol Vis Sci* 2008; **49**: 1571–9.
38. Puche N, Querques G, Benhamou N, et al.. High-resolution spectral domain optical coherence tomography features in adult onset foveomacular vitelliform dystrophy. *Br J Ophthalmol* 2010; **94**: 1190–6.
39. Oishi A, Hata M, Shimoazono M, Mandai M, Nishida A, Kurimoto Y. The significance of external limiting membrane status for visual acuity in age-related macular degeneration. *Am J Ophthalmol* 2010; **150**: 27–32 e21.
40. Spaide RF, Koizumi H, Freund KB. Photoreceptor outer segment abnormalities as a cause of blind spot enlargement in acute zonal occult outer retinopathy-complex diseases. *Am J Ophthalmol* 2008; **146**: 111–20.
41. Miller DT, Kocaoglu OP, Wang Q, Lee S. Adaptive optics and the eye (super resolution OCT). *Eye* 2011; **25**: 321–30.
42. Rowe MP, Engheta N, Easter SS Jr, Pugh EN Jr. Graded-index model of a fish double cone exhibits differential polarization sensitivity. *J Opt Soc Am A Opt Image Sci Vis* 1994; **11**: 55–70.
43. Enoch JM, Hope GM. Directional sensitivity of the foveal and parafoveal retina. *Invest Ophthalmol* 1973; **12**: 497–503.
44. Hoang QV, Linsenmeier RA, Chung CK, Curcio CA. Photoreceptor inner segments in monkey and human retina: mitochondrial density, optics, and regional variation. *Vis Neurosci* 2002; **19**: 395–407.
45. Sidman RL. Histochemical studies on photoreceptor cells. *Ann N Y Acad Sci* 1958; **74**: 182–95.
46. Jonnal RS, Kocaoglu OP, Zawadzki RJ, Lee SH, Werner JS, Miller DT. The cellular origins of the outer retinal bands in optical coherence tomography images. *Invest Ophthalmol Vis Sci* 2014; **55**: 7904–18.
47. Hood DC, Lazow MA, Locke KG, Greenstein VC, Birch DG. The transition zone between healthy and diseased retina in patients with retinitis pigmentosa. *Invest Ophthalmol Vis Sci* 2011; **52**: 101–8.
48. Kitaguchi Y, Kusaka S, Yamaguchi T, Mihashi T, Fujikado T. Detection of photoreceptor disruption by adaptive optics fundus imaging and Fourier-domain optical coherence tomography in eyes with occult macular dystrophy. *Clin Ophthalmol* 2011; **5**: 345–51.
49. Oh J, Smiddy WE, Flynn HW Jr, Gregori G, Lujan B. Photoreceptor inner/outer segment defect imaging by spectral domain OCT and visual prognosis after macular hole surgery. *Invest Ophthalmol Vis Sci* 2010; **51**: 1651–8.
50. Ito N, Kameya S, Gocho K, et al.. Multimodal imaging of a case of peripheral cone dystrophy. *Doc Ophthalmol* 2015; **130**: 241–51.
51. Fritsche LG, Loenhardt T, Janssen A, et al.. Age-related macular degeneration is associated with an unstable ARMS2 (LOC387715) mRNA. *Nat Genet* 2008; **40**: 892–6.
52. Meadway A, Wang X, Curcio CA, Zhang Y. Microstructure of subretinal drusenoid deposits revealed by adaptive optics imaging. *Biomed Opt Express* 2014; **5**: 713–27.
53. Spaide RF. Outer retinal atrophy after regression of subretinal drusenoid deposits as a newly recognized form of late age-related macular degeneration. *Retina* 2013; **33**: 1800–8.

54. Pilotto E, Benetti E, Convento E, *et al.*. Microperimetry, fundus autofluorescence, and retinal layer changes in progressing geographic atrophy. *Can J Ophthalmol* 2013; **48**: 386–93.
55. Sayanagi K, Sharma S, Kaiser PK. Photoreceptor status after antivascular endothelial growth factor therapy in exudative age-related macular degeneration. *Br J Ophthalmol* 2009; **93**: 622–6.
56. Shin HJ, Chung H, Kim HC. Association between foveal microstructure and visual outcome in age-related macular degeneration. *Retina* 2011; **31**: 1627–36.
57. Hayashi H, Yamashiro K, Tsujikawa A, Ota M, Otani A, Yoshimura N. Association between foveal photoreceptor integrity and visual outcome in neovascular age-related macular degeneration. *Am J Ophthalmol* 2009; **148**: 83–9 e81.
58. Nguyen MH, Witkin AJ, Reichel E, *et al.*. Microstructural abnormalities in MEWDS demonstrated by ultrahigh resolution optical coherence tomography. *Retina* 2007; **27**: 414–8.
59. Hashimoto H, Kishi S. Ultra-wide-field fundus autofluorescence in multiple evanescent white dot syndrome. *Am J Ophthalmol* 2015; **159**: 698–706.
60. Lee GE, Lee BW, Rao NA, Fawzi AA. Spectral domain optical coherence tomography and autofluorescence in a case of acute posterior multifocal placoid pigment epitheliopathy mimicking Vogt–Koyanagi–Harada disease: case report and review of literature. *Ocul Immunol Inflamm* 2011; **19**: 42–7.
61. Darugar A, Mathian A, Lehoang P, Bodaghi B. Acute posterior multifocal placoid pigment epitheliopathy as the initial manifestation of sarcoidosis. *J Ophthalmic Vision Res* 2011; **6**: 338–43.
62. Channa R, Ibrahim M, Sepah Y, *et al.*. Characterization of macular lesions in punctate inner choroidopathy with spectral domain optical coherence tomography. *J Ophthalmic Inflamm Infect* 2012; **2**: 113–20.
63. Hua R, Liu L, Chen L. Evaluation of the progression rate of atrophy lesions in punctate inner choroidopathy (PIC) based on autofluorescence analysis. *Photodiagnosis Photodyn Ther* 2014; **11**: 565–9.
64. Zhang X, Zuo C, Li M, Chen H, Huang S, Wen F. Spectral-domain optical coherence tomographic findings at each stage of punctate inner choroidopathy. *Ophthalmology* 2013; **120**: 2678–83.
65. Ohta K, Sato A, Fukui E. Spectral domain optical coherence tomographic findings at convalescent stage of acute zonal occult outer retinopathy. *Clin Ophthalmol* 2009; **3**: 423–8.
66. Nakao S, Kaizu Y, Yoshida S, Iida T, Ishibashi T. Spontaneous remission of acute zonal occult outer retinopathy: follow-up using adaptive optics scanning laser ophthalmoscopy. *Graefes Archive Clin Exp Ophthalmol* 2015; **253**: 839–43.
67. Chai Y, Yamazaki H, Fujinami K, Tsunoda K, Yamamoto S. Case of acute zonal occult outer retinopathy with abnormal pattern visual evoked potentials. *Clin Ophthalmol* 2011; **5**: 1235–41.
68. Ko TH, Fujimoto JG, Duker JS, *et al.*. Comparison of ultrahigh- and standard-resolution optical coherence tomography for imaging macular hole pathology and repair. *Ophthalmology* 2004; **111**: 2033–43.
69. Wakabayashi T, Fujiwara M, Sakaguchi H, Kusaka S, Oshima Y. Foveal microstructure and visual acuity in surgically closed macular holes: spectral-domain optical coherence tomographic analysis. *Ophthalmology* 2010; **117**: 1815–24.
70. Inoue M, Morita S, Watanabe Y, *et al.*. Preoperative inner segment/outer segment junction in spectral-domain optical coherence tomography as a prognostic factor in epiretinal membrane surgery. *Retina* 2011; **31**: 1366–72.
71. Mitamura Y, Hirano K, Baba T, Yamamoto S. Correlation of visual recovery with presence of photoreceptor inner/outer segment junction in optical coherence images after epiretinal membrane surgery. *Br J Ophthalmol* 2009; **93**: 171–5.
72. Oster SF, Mojana F, Brar M, Yuson RM, Cheng L, Freeman WR. Disruption of the photoreceptor inner segment/outer segment layer on spectral domain-optical coherence tomography is a predictor of poor visual acuity in patients with epiretinal membranes. *Retina* 2010; **30**: 713–8.
73. Akkoyun I, Yilmaz G. Optical coherence tomography: anatomic and functional outcome after scleral buckling surgery in macula-off rhegmatogenous retinal detachment. *Klin Monbl Augenheilkd* 2013; **230**: 814–9.
74. Sheth S, Dabir S, Natarajan S, Mhatre A, Labauri N. Spectral domain-optical coherence tomography study of retinas with a normal foveal contour and thickness after retinal detachment surgery. *Retina* 2010; **30**: 724–32.
75. Shimoda Y, Sano M, Hashimoto H, Yokota Y, Kishi S. Restoration of photoreceptor outer segment after vitrectomy for retinal detachment. *Am J Ophthalmol* 2010; **149**: 284–90.
76. Paunescu LA, Ko TH, Duker JS, *et al.*. Idiopathic juxtafoveal retinal telangiectasis: new findings by ultrahigh-resolution optical coherence tomography. *Ophthalmology* 2006; **113**: 48–57.
77. Murakami T, Tsujikawa A, Ohta M, *et al.*. Photoreceptor status after resolved macular edema in branch retinal vein occlusion treated with tissue plasminogen activator. *Am J Ophthalmol* 2007; **143**: 171–3.
78. Shin HJ, Chung H, Kim HC. Association between integrity of foveal photoreceptor layer and visual outcome in retinal vein occlusion. *Acta Ophthalmol* 2011; **89**: e35–40.
79. Matsumoto H, Sato T, Kishi S. Outer nuclear layer thickness at the fovea determines visual outcomes in resolved central serous chorioretinopathy. *Am J Ophthalmol* 2009; **148**: 105–10 e101.
80. Ojima Y, Tsujikawa A, Yamashiro K, Ooto S, Tamura H, Yoshimura N. Restoration of outer segments of foveal photoreceptors after resolution of central serous chorioretinopathy. *Jpn J Ophthalmol* 2010; **54**: 55–60.
81. Maheshwary AS, Oster SF, Yuson RM, Cheng L, Mojana F, Freeman WR. The association between percent disruption of the photoreceptor inner segment-outer segment junction and visual acuity in diabetic macular edema. *Am J Ophthalmol* 2010; **150**: 63–7 e61.
82. Sakamoto A, Nishijima K, Kita M, Oh H, Tsujikawa A, Yoshimura N. Association between foveal photoreceptor status and visual acuity after resolution of diabetic macular edema by pars plana vitrectomy. *Graefes Arch Clin Exp Ophthalmol* 2009; **247**: 1325–30.

83. Shin HJ, Lee SH, Chung H, Kim HC. Association between photoreceptor integrity and visual outcome in diabetic macular edema. *Graefes Arch Clin Exp Ophthalmol* 2012; **250**: 61–70.
84. Scarinci F, Jampol LM, Linsenmeier RA, Fawzi AA. Association of diabetic macular nonperfusion with outer retinal disruption on optical coherence tomography. *JAMA Ophthalmol* 2015; **133**: 1036–44.
85. Shen J, Feng L, Li Y. Ocular syphilis: an alarming infectious eye disease. *Int J Clin Exp Med* 2015; **8**: 7770–7.
86. Yamamoto K, Takahashi H, Kanno M, Noda Y, Fujino Y. Changes in parafoveal retinal thickness and subfoveal choroidal thickness in a patient with dengue fever-associated maculopathy. *J Ophthalmic Inflamm Infect* 2013; **3**: 63.
87. Podoleanu AG, Dobre GM, Webb DJ, Jackson DA. Simultaneous en-face imaging of two layers in the human retina by low-coherence reflectometry. *Opt Lett* 1997; **22**: 1039–41.
88. Wanek J, Zelkha R, Lim JI, Shahidi M. Feasibility of a method for en face imaging of photoreceptor cell integrity. *Am J Ophthalmol* 2011; **152**: 807–14 e801.
89. Nunes RP, Gregori G, Yehoshua Z, et al.. Predicting the progression of geographic atrophy in age-related macular degeneration with SD-OCT en face imaging of the outer retina. *Ophthalmic Surg, Lasers Imaging Retin* 2013; **44**: 344–59.
90. Zhang Y, Rha J, Jonnal R, Miller D. Adaptive optics parallel spectral domain optical coherence tomography for imaging the living retina. *Opt Express* 2005; **13**: 4792–811.
91. Drexler W. Ultrahigh-resolution optical coherence tomography. *J Biomed Opt* 2004; **9**: 47–74.
92. Ting DS, Cheung GC, Lim LS, Yeo IY. Comparison of swept source optical coherence tomography (SS-OCT) and spectral domain optical coherence tomography (SD-OCT) in polypoidal choroidal vasculopathy. *Clin Experiment Ophthalmol* 2015; **43**: 815–9.
93. Imai A, Toriyama Y, Iesato Y, Hirano T, Murata T. En face swept-source optical coherence tomography detecting thinning of inner retinal layers as an indicator of capillary nonperfusion. *Eur J Ophthalmol* 2015; **25**: 153–8.
94. Flores-Moreno I, Arias-Barquet L, Rubio-Caso MJ, Ruiz-Moreno JM, Duker JS, Caminal JM. En face swept-source optical coherence tomography in neovascular age-related macular degeneration. *Br J Ophthalmol* 2015.