

The Importance of Ocular, Systemic and Psychosocial Factors in the Management of the Diabetic Macular Oedema Patient with Anti-vascular Endothelial Growth Factor Therapy

Expert Recommendations and Literature Review

Richard Gale,¹ Angela Carneiro,² Julie de Zaeytijd,³ Paul Dodson,⁴ Sascha Fauser,⁵ João Figueira,⁶ Michael Larsen,⁷ Nicolas Leveziel,⁸ Michael Kapusta,⁹ Jose M Ruiz-Moreno,¹⁰ Enrico Peiretti,¹¹ Christian Pruenste,¹² Reiner Schlingemann¹³ and Christoph Scholda¹⁴

1. York Teaching Hospital NHS Foundation Trust, UK; 2. University of Porto and Department of Ophthalmology of Hospital São João, Portugal; 3. Ghent University Hospital, Belgium; 4. Heartlands Hospital & Aston University, UK; 5. University Hospital of Cologne, Germany; 6. Centro Hospitalar e Universitário de Coimbra and Faculty of Medicine of the University of Coimbra, Portugal; 7. Glostrup Hospital and National Eye Clinic, University of Copenhagen, Denmark; 8. University Hospital Center of Poitiers, France; 9. McGill University, Canada; 10. Castilla La Mancha University, Spain; 11. University of Cagliari, Italy; 12. University of Basel, Switzerland; 13. Academic Medical Center, The Netherlands; 14. Medical University of Vienna, Austria;

A comprehensive list of patient features that influence the management of patients with diabetic macular oedema (DMO) is discussed. These features are grouped into three overarching themes: ocular, systemic and psychosocial. Consensus statements about the relative importance of these features, supported by the literature, were formed by a panel of retinal experts. The major drivers influencing the management of DMO with anti-vascular endothelial growth factor (anti-VEGF) therapy are undoubtedly ocular specific, in particular visual acuity and optical coherence tomography (OCT) central retinal thickness. Systemic factors, such as control of glycated haemoglobin (HbA_{1c}), blood pressure and serum lipid estimations, have limited direct influence on DMO management although they remain important considerations to communicate to the primary diabetic physician. A greater understanding is required on how many other factors, in particular psychosocial factors, influence the care of the DMO patient.

Keywords

Diabetic macular oedema (DMO); management recommendations; anti-vascular endothelial growth factor; visual acuity; OCT

Disclosure: TBC

Acknowledgements: Medical writing assistance was provided by Catherine Amey and James Gilbert, Touch Medical Media UK, funded by Bayer.

Authorship: All named authors meet the International Committee of Medical Journal Editors (ICMJE) criteria for authorship of this manuscript, take responsibility for the integrity of the work as a whole, and have given final approval to the version to be published.

Open Access: This article is published under the Creative Commons Attribution Noncommercial License, which permits any non-commercial use, distribution, adaptation and reproduction provided the original author(s) and source are given appropriate credit.

Received: 2 November 2016

Accepted: 6 December 2016

Citation: *European Ophthalmic Review*, 2016;10(2):xx-xx

Corresponding Author: Richard Gale, York Teaching Hospital NHS Foundation Trust, York, YO31 8HE, UK. E: richard.gale@york.nhs.uk

Support: The publication of this article was supported by Bayer. The views and opinions expressed are those of the author and do not necessarily reflect those of Bayer.

Diabetic macular oedema (DMO) is one of the leading causes of visual impairment in working adults.¹ The implications of blinding due to DMO, including the loss of productivity and reduced quality of life (QoL), lead to a considerable socioeconomic burden on communities.²⁻⁵ Intravitreal treatment options, particularly with anti-vascular endothelial growth factor (anti-VEGF) agents, have shown the potential to reduce visual loss far beyond that achieved with laser therapy alone.⁶⁻¹⁰ Applying principles learnt from key clinical trials to the real world setting requires an understanding of the many patient-related factors that determine when and how treatment is started and continued. It is important for clinicians to not only understand such factors but also the evidence base for such.

The aim of this article is to bring clarity to the available data on the key patient-related factors that influence the treatment of DMO, by reviewing the literature, forming consensus statements and providing recommendations. Discussions are based around the optimum scenario in an ideal world where certain barriers were removed, for example, costing, reimbursement and resource capacity. Due to the broad nature of this topic and because anti-VEGF therapy is the commonest intravitreal intervention for DMO, the scope of the discussion is largely limited to treatment with anti-VEGF therapy.

Methods

A panel of 15 retinal experts (see *Table 1*) met on 14–15 November 2014 in Zurich, Switzerland, where the need for clarity of the factors that influence DMO management was initially discussed. The panel met again on 10–11 April 2015 in Berlin, Germany, and on 6–7 November 2015 in Paris, France, to take the work forward. Based on a Delphi-style methodology,¹¹ a range of factors in the management of a patient with DMO were identified and the evidence for, and relative importance of each, was discussed. A literature search complemented this process. PubMed/Medline and EMBASE searches from 1 January 2005 to 22 June 2016 were carried out using these factors as exploded mesh terms that occurred in the publication title (i.e. the term itself and a series of similar terms and subtopic terms within a particular category as used to classify each record by the database providers). Animal studies and non-English language papers were excluded. Owing to the paucity of material identified through the searches on

Table 1: List of panel members who convened on 14–15 November 2014 in Zurich, Switzerland, to discuss important factors affecting management of patients with diabetic macular oedema

Panel Member	Affiliation
Dr Richard Gale	York Teaching Hospital NHS Foundation Trust, UK
Dr Angela Carneiro	University of Porto and Department of Ophthalmology of Hospital São João, Portugal
Dr Julie de Zaeytijd	Ghent University Hospital, Belgium
Professor Paul Dodson	Heartlands Hospital & Aston University, UK
Dr Sascha Fauser	University Hospital of Cologne, Germany
Dr João Figueira	Centro Hospitalar e Universitário de Coimbra and Faculty of Medicine of the University of Coimbra, Portugal
Dr Gerhard Kieselbach	Medical University Innsbruck, Austria
Professor Michael Larsen	Glostrup Hospital and National Eye Clinic, University of Copenhagen, Denmark
Dr Nicolas Leveziel	University Hospital Center of Poitiers, France
Dr Michael Kapusta	McGill University, Canada
Dr Jose M Ruiz-Moreno	Castilla La Mancha University, Spain
Dr Enrico Peiretti	University of Cagliari, Italy
Professor Christian Prunte	University of Basel, Switzerland
Dr Reiner Schlingemann	Academic Medical Center, The Netherlands
Dr Christoph Scholda	Medical University of Vienna, Austria

Subsequent meetings were held in Berlin, Germany (10–11 April 2015) and in Paris, France (6–7 November 2015).

occupation/profession, co-morbidities, compliance and family history, it was decided to broaden the searches to papers where the specified factor was not used as a major term. After removing duplicates, the search results were screened for relevance by a single retinal expert (Richard Gale). Recommendations were formed and agreed by the panel.

Results

Factors were categorised into three overarching themes: ocular-specific, systemic influences and psychosocial factors (see *Tables 2 and 3*). Database searching identified 547 records and a further 140 records were provided from expert opinion (687 in total). Removing duplicates left 665 results, of which 606 were excluded by expert screening. In total, therefore, 59 full text articles were accessed and relevant articles discussed.

Ocular-specific factors that influence the management of diabetic macular oedema with anti-vascular endothelial growth factor therapy (Recommendation 1)

Many overlapping definitions of DMO are currently used, for example, clinically significant macular oedema (CSMO), centre involving vision affecting (CIVA) oedema and focal or diffuse oedema.^{12,13} Using the correct definition is central to aligning treatment with the key clinical trials.

In 1985, the Early Treatment Diabetic Retinopathy Study (ETDRS) group defined CSMO,¹² based on clinical examination alone: presence of any retinal thickening at or within 500 µm of the foveal centre, lipid exudates at or within 500 µm of the foveal centre with adjacent thickening or an area of retinal thickening at one or more Macular Photocoagulation Study DA (1 disk area \cong 1.767 mm²) within one disk diameter (1.5 mm) of the foveal centre. This definition remains useful in patients undergoing macular laser therapy; however, it is not the driver when anti-VEGF therapy is used. The inclusion criteria for the majority of the pivotal trials uses a form of 'centre involving, vision affecting' criteria to initiate treatment. *Tables 4 and 5* demonstrate that inclusion criteria for the RISE and RIDE, Protocol I and T and DA VINCI studies required a combination of vision

Table 2: Three overarching themes of identified factors that affect management of patients with diabetic macular oedema

Ocular	Systemic	Psychosocial
DMO definitions, e.g. Clinically significant macular oedema	Glycated haemoglobin	Profession
Imaging Optical coherence tomography Fundus fluorescein angiography	Blood pressure	Age
Visual acuity	Lipids	Compliance
Ocular co-morbidities, e.g. Cataract Glaucoma	Thiazolidinedione treatment	
Unilateral and bilateral treatment	Body mass index	Quality of life Patient-reported symptoms
	Pregnancy	
	Stroke or myocardial infarction within 3 months	

loss and documented retinal thickening.^{6,7,12} It is worth noting, however, the inclusion criteria for the RESTORE study did not require a specific CRT measurement but a clinical diagnosis of diffuse or focal thickening.⁹ Focal DMO was defined as more than 67% of leakage originated from leaking microaneurysms (MAs) in the whole oedema area or 30–67% leakage from MAs in the whole oedema area, but >67% of the leakage originated from MAs in the central subfield.⁹ Diffuse DMO was defined as <33% of leakage from MAs, the rest from diffuse leaking capillaries in the whole oedema area or 30–67% leakage from MAs, but <33% of the leakage originated from MAs in the central subfield. This definition is somewhat convoluted and is not clinically important as a sub-analysis of the RESTORE data demonstrated neither group performed better than the other.⁹

Table 3: Recommendations on the importance of ocular, systemic and psychosocial factors on the management of the patient with diabetic macular oedema

Number	Recommendation
A. Ocular-specific	
Definition	
1	Use of 'centre involving vision affecting DMO' is the most useful term in describing DMO to be treated with anti-VEGF therapy and is the primary driver to commence therapy
Optical coherence tomography	
2	In conjunction with clinical context, OCT is the principal tool used to diagnose DMO. Clinical trial evidence provides support for its use to guide retreatment. The central 1 mm ² subfield thickness is the most useful measurement
Fundus fluorescein angiography	
3	While FFA continues to have a role in helping excluded differential diagnosis and documenting macular ischaemia it otherwise has little place in guiding commencement of therapy or retreatment
Visual acuity	
4	VA is one of the principle factors used to guide commencement and retreatment of therapy, it is a proxy measure of QoL. Caution is required when VA is used to guide treatment because of its natural variability in repeat measurements between visits
5	Patients with unilateral or bilateral DMO should be treated the same although compromise in the treatment regime may have to be made for logistical reasons
Co-morbidities	
6	The influence of ocular co-morbidities on the potential of therapy to restore vision need to be considered
B. Systemic influences	
Glycated haemoglobin	
7	HbA _{1c} has no influence on the decision to commence therapy or the decision to retreat. Communication with the physician managing the general diabetes is important to optimise control, which in turn optimises the retinopathy in the long term
Blood pressure	
8	BP, unless very high, has little influence on the decision to commence therapy or the decision to retreat. Communication with the physician managing the general diabetes is important to optimise control, which in turn optimises the retinopathy in the long term
Lipids	
9	Serum lipid status has no influence on the decision to commence therapy or the decision to retreat. Communication with the physician managing the general diabetes is important to optimise control, which in turn optimises the retinopathy in the long term
TZDs	
10	TZDs should be withdrawn by the prescribing doctor if there is co-existing DMO. It is reasonable to wait a period several weeks to months to see the effect of this before commencing therapy.
Body mass index	
11	BMI has no influence on whether a patient should be treated with an anti-VEGF agent. BMI optimisation via the patient's diabetic physician is recommended.
Pregnancy	
12	It is imperative that treating physicians routinely ask for and if necessary test for pregnancy as this remains a contraindication for anti-VEGF therapy
Stroke and myocardial infarction	
13	Recent stroke or myocardial infarction (<3 months) remains a contraindication for anti-VEGF therapy. Treatment within these time frames requires discussion with the patient about the relative risk and benefits of doing so
C. Psychosocial factors	
QoL	
14	Improving QoL utilities and symptoms of DMO are the ultimate goal in the management of DMO. They are important considerations but may be too crude or imprecise to specifically guide treatment
Age	
15	In the adult population there is no evidence that age, per se, should influence treatment with anti-VEGF agent
Profession	
16	A patient's profession may influence their visual requirements (QoL influences) and compliance with treatment and so should be discussed
Compliance	
17	The importance of compliance should be discussed prior to commencement of therapy. Non-compliant patients have not been specifically studied, however, the result of poor attendance is likely to be compromised treatment benefit.

BMI = body mass index; BP = blood pressure; CV = cardiovascular; DMO = diabetic macular oedema; FFA = fundus fluorescein angiography; HbA_{1c} = glycated haemoglobin; QoL = quality of life; OCT = optical coherence tomography; anti-VEGF = anti-vascular endothelial growth factor agents.

Table 4: Key inclusion/exclusion criteria in RESTORE,⁹ RISE/RIDE,¹⁵ DRCR.net Protocol I,⁸ Protocol T,¹⁰ DA VINCI⁷ and VISTA/VIVID¹⁶

Key Factor	Study					
	Restore ⁹	RISE/RIDE ¹⁵	DRCR.net Protocol I ⁸	DRCR.net Protocol T ¹⁰	DA VINCI ⁷	VISTA/VIVID ¹⁶
OCT	–	CRT ≥275 µm (inclusion)	405 µm on time-domain OCT (inclusion)	OCT ≥250 µm on Zeiss Stratus; ≥320 if male or ≥305 if female on Heidelberg Spectralis; ≥305 if male or ≥290 if female on Zeiss Cirrus	CRT ≥250 µm (inclusion)	Retinal thickness ≥300 µm as assessed by OCT (inclusion)
VA	BCVA between 78 and 39 inclusive (inclusion)	BCVA 20/40 to 20/320 Snellen equivalent (inclusion)	20/32 or worse (inclusion)	≤78 (20/32 or worse) and ≥24 (20/320 or better) (inclusion)	BCVA letter score 73 to 24 (20/40–20/320) (inclusion)	BCVA letter score 73–24 (20/40–20/320) (inclusion)
HbA1c	≤10% (inclusion)	>12% (exclusion)	Unsuitable glycaemic control as deemed by the investigator (exclusion)	–	–	Uncontrolled diabetes mellitus in the opinion of investigator (VISTA) or as defined by HbA _{1c} >12% (VIVID) (exclusion)
BP	>160 mmHg SBP or >100 mg Hg DBP (exclusion)	Uncontrolled hypertension (exclusion)	BP >180/110 (SBP >180 OR DBP >110) (exclusion)	Blood pressure >180/110 (systolic >180 OR diastolic >110) (exclusion)	SBP > 180 mmHg or >160 mmHg on 2 consecutive measurements; DBP >100 mmHg (exclusion)	SBP >160 mmHg or DBP >95 mmHg while sitting (exclusion)
Lipids	–	–	–	–	–	–
Age	≥18 years (inclusion)	≥18 years (inclusion)	≥18 years (inclusion)	≥18 years (inclusion)	≥18 years (inclusion)	≥18 years (inclusion)

BP = blood pressure; CRT = central retinal thickness; DBP = diastolic blood pressure DRCR.net = Diabetic Retinopathy Clinical Research Network; HbA_{1c} = glycated haemoglobin; OCT = optical coherence tomography; SBP = systolic blood pressure; VA = visual acuity.

Optical coherence tomography findings (Recommendation 2)

Optical coherence tomography (OCT) is the most widely used tool to diagnose and manage DMO. The central 1 mm subfield is the most useful parameter in determining initiation and retreatment. Patients were recruited into the DRCR.net Protocol I who had DMO involving the fovea that, amongst other criteria, required a mean OCT central subfield thickness on OCT >250 µm on time-domain OCT.¹⁴ Inclusion for participation in DRCR.net Protocol T included: central subfield thickness on OCT ≥250 µm on Zeiss Stratus; ≥320 if male or ≥305 if female on Heidelberg Spectralis; ≥305 if male or ≥290 if female on Zeiss Cirrus and definite retinal thickening on clinical exam due to DMO involving the centre of the macula.⁶ In the RISE and RIDE trials, eligible participants required macular oedema with a central subfield thickness ≥275 µm on time-domain OCT.¹⁵ In the DA VINCI study, clinically significant DMO with centre involvement of the fovea was defined as a central subfield measurement of ≥250 µm on time-domain OCT.⁷ Central DMO involvement was defined as retinal thickening involving 1 mm central (OCT) subfield thickness (CST) in the VISTA and VIVID trials.¹⁶

Structural OCT does not provide dynamic information about macular perfusion. OCT angiography (OCT-A) is a developing technology and, as yet, its role in the routine assessment of the DMO patient is still to be fully defined. Used in parallel with spectral domain (SD OCT), OCT-A presents the opportunity to learn more about the disease.^{17,18}

Fundus fluorescein angiography findings (Recommendation 3)

Fundus fluorescein angiography (FFA) is no longer commonly used to make the diagnosis of DMO. On occasions, FFA is important to differentiate DMO from other vascular disorders, such as choroidal neovascularisation, retinal vein occlusion or macular teleangiectasia. There is little direct

evidence that FFA has a role in the management of DMO with anti-VEGF therapy. The key clinical trials did not specify FFA features as a part of inclusion criteria for treatment nor for retreatment (RESTORE, RISE and RIDE, Protocol I and T, DA VINCI). FFA is useful in assessing the degree of enlargement and irregularity of the foveal avascular zone, i.e. macular ischaemia. However, patients with macular ischaemia were studied in a subgroup analysis in the RESTORE study and no difference was found in their outcome – implying patients with or without macular ischaemia should be treated similarly.⁹ A caveat is that patients were excluded from these studies if there was a reason why vision may not improve and so patients with very severe macular ischaemia may not have been enrolled.

Visual acuity (Recommendation 4)

Visual acuity (VA) (ETDRS score) is a key factor influencing commencement of therapy and retreatment (see *Tables 4* and *5*). Indeed VA assessment was used as an inclusion criteria in RESTORE,⁹ DRCR.net Protocols I⁸ and T,¹⁰ DA VINCI⁷ and VISTA and VIVID studies¹⁶ and furthermore to help guide retreatment (see *Tables 4* and *5*).^{8–10,16} The key clinical trials studying the use of anti-VEGF therapy in DMO only included patients with reduced vision so currently there is no evidence that supports the benefit of anti-VEGF therapy in those with unaffected, very good or very poor VA.

The DRCR.net published findings of a refracted ETDRS coefficient of repeatability of 5–13 letters between visits (the worse the VA the higher the coefficient). Caution is therefore suggested when change in VA is used prospectively as a determination of a true measure of change in visual function.

Unilateral or bilateral treatment (Recommendation 5)

There is no evidence from the pivotal clinical trials that unilateral or bilateral treatment influences the commencement or retreatment of

Table 5: Retreatment criteria from RESTORE,⁹ RISE/RIDE,¹⁵ DRCR.net Protocol I,⁸ Protocol T,¹⁰ DA VINCI⁷ and VISTA/VIVID¹⁶

Study					
RESTORE ⁹	RISE/RIDE ¹⁵	DRCR.net Protocol I ⁸	DRCR.net Protocol T ¹⁰	DA VINCI ⁷	VISTA/VIVID ¹⁶
1 injection per month continued unless stable VA Treatment suspended if: (1) the investigator's opinion was that no (further) BCVA improvement was attributable to treatment with intravitreal injection at the last 2 consecutive visits, or (2) BCVA letter score ≥ 84 (approximate Snellen equivalent 20/20) was observed at the last 2 consecutive visits	Monthly injections	If VA is ≥ 84 (20/20 or better) or the OCT central subfield thickness is $< 250 \mu\text{m}$, the decision to reinject is at investigator discretion. If VAs < 84 (worse than 20/20) and OCT central subfield thickness $\geq 250 \mu\text{m}$, an injection (or sham) is given except in the case of prior AEs	Every 4 weeks unless VA was 20/20 or better with a central subfield thickness below eligibility criteria	VEGF Trap-Eye 0.5 mg every 4 weeks (0.5q4); 2 mg every 4 weeks (2q4); 2 mg every 8 weeks after 3 initial monthly doses (2q8); or 2 mg dosing as needed after 3 initial monthly doses (2 PRN), or macular laser photocoagulation	Assessed for laser retreatment beginning at week 12. If any ETDRS-defined, clinically significant macular oedema, for which laser has been shown to be beneficial, study eyes in the 2q4 and 2q8 groups received sham laser and those in the laser group received active laser, but not more frequently than every 12 weeks

DRCR.net = Diabetic Retinopathy Clinical Research Network; BCVA = best corrected visual acuity; ETDRS = Early Treatment Diabetic Retinopathy Study; OCT = optical coherence tomography; PRN = pro re nata; VA = visual acuity; VEGF = vascular endothelial growth factor.

patients (RESTORE, RIDE, RISE, Protocol I and T, DA VINCI and VISTA and VIVID studies). Similarly, there is no evidence that a better or worse seeing eye is treated differently. A compromise in treatment regime may have to be made for those receiving bilateral treatment due to logistical reasons.

Ocular co-morbidities (Recommendation 6)

Should a patient have an ocular co-morbidity, physicians need to consider the potential to restrict the effect of any treatment or cause more profound loss. An example may be co-existing foveal-involving atrophy due to age-related macular degeneration. Key studies excluded patients with significant co-morbidities from being enrolled into clinical trials.^{8,9,15,19} VA measurements therefore may not reflect change in DMO due to co-morbidities.

Systemic factors that influence the management of diabetic macular oedema with anti-vascular endothelial growth factor therapy

Blood Glucose Control (Recommendation 7)

The evidence that management with anti-VEGF therapy for DMO should differ based upon their glycated haemoglobin (HbA_{1c}) is weak. An example of this is a post hoc analysis of the RIDE and RISE trials, in which the treatment benefit of ranibizumab for patients with DMO seemed independent of baseline HbA_{1c} .²⁰ A retrospective consecutive case series of 124 patients with DMO treated with anti-VEGF, however, did find that glucose regulation can impact on the response to anti-VEGF therapy.²¹ Patients with improved glucose control during the study had a significantly lower retinal thickness than patients who had a stable or worsening HbA_{1c} (mean final CST of 324.3 versus 390.0 μm , respectively; $p=0.042$). Many studies have excluded patients with a very high HbA_{1c} , including the RESTORE,⁹ RISE/RIDE,¹⁵ DRCR.net Protocols I⁸ and T¹⁰, the DA VINCI⁷ and VISTA and VIVID studies¹⁶ (see Table 4) and so the outcome in this group is not as well known.

Advising and reinforcing the importance of good glucose control at the point of contact during DMO management is good practice for the holistic care of patients with diabetes.²¹⁻²³ Intensive glycaemic control decreases the rate of development and progression of diabetic retinopathy (DR) in type 1 and 2 diabetic mellitus patients. In the primary prevention cohort of the Diabetes Control and Complications Trial (DCCT), there was a 76%

reduction in the adjusted mean risk for retinopathy development for patients who received intensive therapy. In the secondary intervention cohort, the progression of retinopathy was reduced by 54%, and the development of proliferative or severe non-proliferative retinopathy was reduced by 47%.²⁴ The National institute for Health and Excellence (NICE) has suggested a target HbA_{1c} for patients with diabetes of 48 mmol/mol (6.5%).^{25,26} It is also good practice to ensure appropriate arrangements are in place, with either a diabetologist or primary care physician for general diabetic care.

Blood pressure control (Recommendation 8)

There is little evidence that patients should be treated differently based upon their starting BP. The only potential exception is in those with very high BP who were excluded from the major clinical trials and therefore it is not known if their outcome differs (see Table 4).^{7-10,15} Likewise, once treatment has been commenced there is no evidence, based upon these studies, that retreatment should differ depending upon BP.

However, theoretically, higher BP will be associated with an increase in retinal capillary hydrostatic pressure due to the dysfunction of vascular autoregulation in the retina of patients with DR. According to Starling's rules this will augment macular oedema, and indeed there is clinical and experimental evidence that high BP is an independent direct and short-term driver of DMO.²⁷⁻²⁹

In addition, there is solid evidence to support the benefit of tightly controlled blood pressure (BP) on overall retinopathy status. The UKPDS investigated the role of tight BP control on various end points, including retinopathy and visual loss. In the group with tight BP control, a 35% reduction in the progression of retinopathy was observed compared with the group assigned to less tight control. At 9-year follow-up, the group with tight BP control had a 47% reduction in risk of loss of three or more lines of vision (using ETDRS chart).²⁹ Gallego et al. identified that an increase of 10 mm/Hg in systolic BP was associated with a 3-20% increased risk of retinopathy and an increase of 10 mm/Hg in diastolic BP increased risk by 2-30%.³⁰ As such, and based on clinical observations and the theoretical issues raised above, it is good practice to ensure BP is being addressed by the appropriate care physician and to recommend BP control in line with recommendation from organisations such as

NICE and Diabetes UK (BP target: 130/80 mmHg if retinopathy, renal or cerebrovascular damage).²⁵

Serum lipid control (Recommendation 9)

Based upon the pivotal anti-VEGF DMO treatment studies, there is no evidence that serum lipid status should alter the way in which anti-VEGF therapy is given. Neither inclusion criteria for the commencement of therapy or retreatment strategies were guided by serum lipid levels.

Although some study findings suggest that the management of serum lipids may reduce retinopathy progression and the need for treatment,^{31,32} other clinical data have disputed the association between lipids and the severity of DR or DMO. In an ETDRS report, high total cholesterol and low-density lipoprotein (LDL) levels were associated with retinal hard exudates.³² The Multi-Ethnic Study of Atherosclerosis and the Australian Diabetes, Obesity, and Lifestyle Study observed no association between serum lipids and DR.^{33,34} In another study, the level of total cholesterol, triglyceride, high-density lipoprotein (HDL), LDL and very low density lipoprotein (VLDL) was not significantly associated with the severity of DR or existence of macular oedema.³⁵ Furthermore Chew et al. observed that DMO was not significantly associated with high triglyceride levels, total cholesterol or LDL levels when lipid profile was stratified.³⁶

Evidence from the Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) and The Action to Control Cardiovascular Risk in Diabetes (ACCORD) Eye studies have shown that fenofibrate may influence DR. In the FIELD study, fenofibrate (200 mg/day) reduced the need for laser therapy and prevented disease progression in patients with pre-existing DR.^{37,38} A systematic review and meta-analysis on dyslipidemia and DMO was inconclusive in its findings, indicating that further investigation into the relationship between lipid levels and DMO is needed.³⁹

Even if controversy exists about the role of serum lipids for DMO specifically, it is important to provide advice on its control for holistic patient benefit. The current NICE guidelines for lipid modification have recommended that a full lipid profile assessment for people with diabetes should include measurement of total cholesterol, HDL cholesterol, non-HDL cholesterol and triglyceride concentrations.⁴⁰ Diabetes UK recommends total cholesterol level <4.0 mmol/l, LDL levels <2.0 mmol/l, HDL levels ≥1.0 mmol/l in men and ≥1.2 mmol/l in women and triglyceride levels ≤1.7 mmol/l.⁴¹

Co-existing thiazolidinedione medication (Recommendation 10)

Peripheral oedema and fluid retention associated with thiazolidinediones (TZDs) may increase the risk of DMO,⁴² however, the exact mechanism underlying this risk remains unclear but may be related to an independent or secondary effect from related systemic fluid retention, possibly by causing increased retinal capillary hydrostatic pressure and oedema formation by Starling's rules. It is advisable to suspend TZD administration when macula oedema develops, since this action is likely to improve the condition. This should be done as a recommendation to the prescribing physician so alternative medication or strategies can be sought. It remains an option to observe whether DMO persists following suspension of a TZD and whether anti-VEGF therapy need be commenced.

Body mass index (Recommendation 11)

Some studies have demonstrated a relationship between obesity or higher body mass index (BMI) and an increased risk of DR.⁴³⁻⁴⁸ Kaštelan et al. suggested that BMI, in conjunction with poor glycaemic control,

hypertension and dyslipidaemia, was associated with DR progression in patients with type 2 diabetes.⁴³ Dirani et al. suggested that patients with diabetes with higher BMI and larger neck circumference were more likely to have DR and more severe stages of DR.⁴⁵ NICE and the UK National Health Service (NHS) Choices guidelines advise that BMI should be maintained within the range 18.5 to 24.9 and that BMI values above 25 increase the risk of type 2 diabetes.^{49,40} There is, however, no evidence available suggesting that, in the context of anti-VEGF therapy, decreasing BMI improves response of DMO, or if it has an influence of how treatment should be administered.

Pregnancy or potential pregnancy (Recommendation 12)

Current evidence on the use of anti-VEGF therapy in pregnancy cases is very clear. Given the potential for teratogenicity, pregnancy is a contraindication for anti-VEGF therapy. There have been case reports, however, of unaffected babies born to mothers undergoing anti-VEGF therapy.⁵¹⁻⁵⁴ Clear advice needs to be given on pregnancy prevention during anti-VEGF therapy or for 3 months thereafter. In addition, consideration needs to be given as to the appropriateness of within clinic pregnancy testing.

Stroke and myocardial infarction (Recommendation 13)

There continues to be discussion in the literature about the potential signal of increased risk of cerebrovascular or cardiovascular events with the use of anti-VEGF therapy. Studies do not enrol patients if these events have occurred recently (<3 months). The risk of recurrence of stroke and myocardial infarction when they have recently occurred (<3 months) in the context of anti-VEGF therapy has not been specifically studied and so careful discussion with the patient about the potential risk and benefits of such treatment is required in this circumstance.^{7,9,10,15,55}

Psychosocial factors that influence the management of diabetic macular oedema with anti-vascular endothelial growth factor therapy

There is little good quality evidence that psychosocial factors influence the use of anti-VEGF therapy in the management of DMO; therefore, the statements in this section are based on opinion.

Quality of life and patient-reported symptoms (Recommendation 14)

QoL utilities and patient-reported symptoms were not used as selection or retreatment criteria in the RESTORE,⁹ RISE/RIDE,¹⁵ DRCR.net Protocols I¹⁴ and T¹⁰ or VISTA and VIVID studies.¹⁶

Age (Recommendation 15)

There is no high-level evidence supporting any association between the influence of adult age on the response to anti-VEGF therapy or alteration in how treatment should be administered. It is noteworthy that the age limitation in the key studies was ≥18 years, without an upper limit (see *Table 4*).^{9,10,15,19,55}

Profession (Recommendation 16)

There are data demonstrating that a patient's working profession is restricted and their employment is made more expensive due to visual impairment by DR.^{2,56-58} There are, however, no data to suggest that a patients' profession has any influence on DMO diagnosis or management.

Compliance (Recommendation 17)

Anti-VEGF therapy in non-compliance patients was not studied in any of the work identified by the literature search. Furthermore, trial participants are not usually included in key clinical trials if the recruiting physicians feel that they will not be compliant with study visits.

People with both vision-threatening DR and psychosocial problems may have significantly reduced levels of functioning compared with psychologically healthy counterparts.⁵⁹ This may have a negative impact on compliance, diabetes control and subsequent outcome. Pivotal studies have not specifically studied this group.

Conclusions

The evidence for the major drivers influencing the management of DMO with anti-VEGF therapy are undoubtedly ocular specific: VA and OCT findings. The panel discussed how patients may decide not to commence therapy though, even in the presence of reduced acuity and a thickened OCT macular scan. This is not an uncommon scenario particularly if a patient was asymptomatic or does not consider the symptoms to be sufficiently troublesome. The evidence for not commencing therapy, or indeed deferring therapy with or without macular laser for a significant period of time (74 weeks or more), demonstrates that VA outcomes are inferior compared with those commencing therapy early.⁶⁰

Although there is little direct evidence that control of systemic factors such as HBA_{1c}, BP and dyslipidemia influence how therapy should be given, there are signals that in the short- and long-term (HBA_{1c} and dyslipidemia) controlling these factors can help with the control of DMO and DR. It is important, not only from this point of view, but also for the holistic care of patients to recommend optimal systemic risk

factor control. This should be achieved through communication with the physician in charge of the blood glucose management whether they are a primary care physician or diabetologist.

Psychosocial factors are important in influencing how patients are managed. Ultimately it is symptoms and QoL that physicians are trying to improve and indeed often are guided by these to influence commencement of treatment. As symptoms and QoL utilities can be either imprecise or relatively crude measures, they are not used as primary treatment endpoints in clinical trials and so no evidence exists to support their use as specific treatment endpoints. Visual function QoL has been shown to improve in key studies though, although EuroQoL five dimensions questionnaire (EQ-5D) did not improve in the RESTORE study. The means by which psychosocial factors influence how patients are managed is so far poorly studied.

There are a number of potential factors that influence the way in which the DMO patient is managed, but many lack a firm evidence base. This lack of evidence raises the risk that treatment decisions may be partly based on assumptions. Better and more extensive evidence to support DMO treatment remains an unmet need. Physicians treating DMO are therefore pointed towards a list of key ocular-specific, systemic and psychosocial considerations that are vital in deciding when to commence or retreat a patient with DMO (see *Table 2*) and these may help improve outcomes. □

- Klein R, Knudtson MD, Lee KE, et al., The Wisconsin Epidemiologic Study of Diabetic Retinopathy XXIII: the twenty-five-year incidence of macular edema in persons with type 1 diabetes, *Ophthalmology*, 2009;116:497–503.
- Brook RA, Kleinman NL, Patel S, et al., United States comparative costs and absenteeism of diabetic ophthalmic conditions, *Postgrad Me*, 2015;127:455–62.
- Chen E, Looman M, Laouri M, et al., Burden of illness of diabetic macular edema: literature review, *Curr Med Res Opin*, 2010;26:1587–97.
- Gonder JR, Walker VM, Barbeau M, et al., Costs and Quality of Life in Diabetic Macular Edema: Canadian Burden of Diabetic Macular Edema Observational Study (C-REALITY), *J Ophthalmol*, 2014;2014:939315.
- Minassian DC, Owens DR, Reidy A, Prevalence of diabetic macular oedema and related health and social care resource use in England, *Br J Ophthalmol*, 2012;96:345–9.
- van't Veer LJ, Dai H, van de Vijver MJ, et al., Gene expression profiling predicts clinical outcome of breast cancer, *Nature*, 2002;415:530–6.
- Do DV, Schmidt-Erfurth U, Gonzalez VH, et al., The DA VINCI Study: phase 2 primary results of VEGF Trap-Eye in patients with diabetic macular edema, *Ophthalmology*, 2011;118:1819–26.
- Elman MJ, Qin H, Aiello LP, et al., Intravitreal ranibizumab for diabetic macular edema with prompt versus deferred laser treatment: three-year randomized trial results, *Ophthalmology*, 2012;119:2312–8.
- Mitchell P, Bandello F, Schmidt-Erfurth U, et al., The RESTORE study: ranibizumab monotherapy or combined with laser versus laser monotherapy for diabetic macular edema, *Ophthalmology*, 2011;118:615–25.
- Wells JA, Glassman AR, Ayala AR, et al., Aflibercept, bevacizumab, or ranibizumab for diabetic macular edema, *New Engl J Med*, 2015;372:1193–203.
- Brown BB, Delphi Process: A Methodology Used for the Elicitation of Opinions of Experts 1968. Available at: <https://www.rand.org/content/dam/rand/pubs/papers/2006/P3925.pdf> (accessed 12 December 2016).
- Photocoagulation for diabetic macular edema. Early Treatment Diabetic Retinopathy Study report number 1. Early Treatment Diabetic Retinopathy Study research group, *Arch Ophthalmol*, 1985;103:1796–806.
- Bandello F, Midena E, Menchini L, Lanzetta P, Recommendations for the appropriate management of diabetic macular edema: Light on DME survey and consensus document by an expert panel, *Eur J Ophthalmol*, 2016;26:252–61.
- Elman MJ, Aiello LP, Beck RW, et al., Randomized trial evaluating ranibizumab plus prompt or deferred laser or triamcinolone plus prompt laser for diabetic macular edema, *Ophthalmology*, 2010;117:1064–77.e35.
- Nguyen QD, Brown DM, Marcus DM, et al., Ranibizumab for diabetic macular edema: results from 2 phase III randomized trials: RISE and RIDE, *Ophthalmology*, 2012;119:789–801.
- Korobelnik JF, Do DV, Schmidt-Erfurth U, et al., Intravitreal aflibercept for diabetic macular edema, *Ophthalmology*, 2014;121:2247–54.
- Coscas GJ, Lupidi M, Coscas F, et al., Optical coherence tomography angiography versus traditional multimodal imaging in assessing the activity of exudative age-related macular degeneration: A new diagnostic challenge, *Retina*, 2015;35:2219–28.
- Jia Y, Bailey ST, Hwang TS, et al., Quantitative optical coherence tomography angiography of vascular abnormalities in the living human eye, *Proc Natl Acad Sci U S A*, 2015;112:E2395–402.
- Do DV, Nguyen QD, Boyer D, et al., One-year outcomes of the da Vinci Study of VEGF Trap-Eye in eyes with diabetic macular edema, *Ophthalmology*, 2012;119:1658–65.
- Bansal AS, Khurana RN, Wieland MR, et al., Influence of glycosylated hemoglobin on the efficacy of ranibizumab for diabetic macular edema: A post hoc analysis of the RIDE/RISE trials, *Ophthalmology*, 2015;122:1573–9.
- Matsuda S, Tam T, Singh RP, et al., The impact of metabolic parameters on clinical response to VEGF inhibitors for diabetic macular edema, *J Diabetes Complications*, 2014;28:166–70.
- Anon. The relationship of glycemic exposure (HbA_{1c}) to the risk of development and progression of retinopathy in the diabetes control and complications trial, *Diabetes*, 1995;44:968–83.
- Anon. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). UK Prospective Diabetes Study (UKPDS) Group, *Lancet*, 1998;352:837–53.
- Anon. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. The Diabetes Control and Complications Trial Research Group, *N Engl J Med*, 1993;329:977–86.
- National Institute for Health and Care Excellence (NICE) guidelines [ng28] Type 2 diabetes in adults: management. Blood glucose and target levels 2015. Available at: <https://www.nice.org.uk/guidance/ng28/ifp/chapter/blood-glucose-and-target-level> (accessed 7 December 2016).
- National Institute for Health and Care Excellence (NICE) guidelines [NG17] Type 1 diabetes in adults: diagnosis and management. Recommendations 2015. Available at: <https://www.nice.org.uk/guidance/ng17/chapter/1-recommendations> (accessed 7 December 2016).
- Klaassen I, Van Noorden CJ, Schlingemann RO, Molecular basis of the inner blood-retinal barrier and its breakdown in diabetic macular edema and other pathological conditions, *Prog Retin Eye Res*, 2013;34:19–48.
- Matthews DR, Stratton IM, Aldington SJ, et al., Risks of progression of retinopathy and vision loss related to tight blood pressure control in type 2 diabetes mellitus: UKPDS 69, *Arch Ophthalmol*, 2004;122:1631–40.
- Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. UK Prospective Diabetes Study Group, *BMJ*, 1998;317:703–13.
- Galleo PH, Craig ME, Hing S, Donaghue KC, Role of blood pressure in development of early retinopathy in adolescents with type 1 diabetes: prospective cohort study, *BMJ*, 2008;337:a918.
- Lyons TJ, Jenkins AJ, Zheng D, et al., Diabetic retinopathy and serum lipoprotein subclasses in the DCCT/EDIC cohort, *Investigative Ophthalmology & Visual Science*, 2004;45:910–8.
- Lopes-Virella MF, Baker NL, Hunt KJ, et al., High concentrations of AGE-LDL and oxidized LDL in circulating immune complexes are associated with progression of retinopathy in type 1 diabetes, *Diabetes Care*, 2012;35:1333BP40.
- Tapp RJ, Shaw JE, Harper CA, et al., The prevalence of and factors associated with diabetic retinopathy in the Australian population, *Diabetes Care*, 2003;26:1731BP7.
- Wong TY, Klein R, Islam FM, et al., Diabetic retinopathy in a multi-ethnic cohort in the United States, *Am J Ophthalmol*, 2006;141:446BP55.
- Cetin EN, Bulgu Y, Ozdemir S, et al., Association of serum lipid levels with diabetic retinopathy, *Int J Ophthalmol*, 2013;6:346BP9.
- Chew EY, Klein ML, Ferris FL, 3rd, et al., Association of elevated serum lipid levels with retinal hard exudate in diabetic retinopathy. Early Treatment Diabetic Retinopathy Study (ETDRS) Report 22, *Arch Ophthalmol*, 1996;114:1079BP84.
- Chew EY, Davis MD, Danis RP, et al., The effects of medical management on the progression of diabetic retinopathy in persons with type 2 diabetes: the Action to Control Cardiovascular Risk in Diabetes (ACCORD) Eye Study, *Ophthalmology*, 2014;121:2443BP51.
- Keetch AC, Mitchell P, Summanen PA, et al., Effect of fenofibrate on the need for laser treatment for diabetic retinopathy (FIELD study): a randomised controlled trial, *Lancet*, 2007;370:1687BP97.
- Das R, Kerr R, Chakravarthy U, Hogg RE, Dyslipidemia and diabetic macular edema: A systematic review and meta-analysis, *Ophthalmology*, 2015;122:1820BP7.
- Landfeldt E, Lindgren P, Bell CF, et al., The burden of Duchenne muscular dystrophy: An international, cross-sectional study, *Neurology*, 2014;83:529–36.
- Mendell JR, Shilling C, Leslie ND, et al., Evidence-based path to newborn screening for Duchenne muscular dystrophy, *Ann Neurol*, 2012;71:304BP13.
- Ryan EH, Jr., Han DP, Ramsay RC, et al., Diabetic macular edema associated with glitazone use, *Retina*, 2006;26:562BP70.
- Kaestel S, Tomic M, Gverovic Antunica A, et al., Body mass index: a risk factor for retinopathy in type 2 diabetic patients, *Mediators Inflamm*, 2013;2013:436329.
- Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). UK Prospective Diabetes Study (UKPDS) Group, *Lancet*, 1998;352:854BP65.
- Dirani M, Xie J, Fenwick E, et al., Are obesity and anthropometry risk factors for diabetic retinopathy? The diabetes management project, *Investigative Ophthalmology & Visual Science*, 2011;52:4416BP21.
- Li X, Wang Z, Prevalence and incidence of retinopathy in elderly diabetic patients receiving early diagnosis and treatment, *Exp Ther Med*, 2013;5:1393BP6.
- Klein R, Klein BE, Moss SE, Visual impairment in diabetes, *Ophthalmology*, 1984;91:1BP9.
- Klein R, Klein BE, Moss SE, Is obesity related to microvascular and macrovascular complications in diabetes? The Wisconsin Epidemiologic Study of Diabetic Retinopathy, *Arch Int Med*, 1997;157:650BP6.

49. Obesity: identification, assessment and management BP NICE guidelines [cg189]. 2014.
50. Reduce your diabetes risk. 2015.
51. Polizzi S, Mahajan VB, Intravitreal anti-VEGF injections in pregnancy: Case series and review of literature, *J Ocul Pharmacol Ther*, 2015;31:605BP10.
52. Polizzi S, Ferrara G, Restaino S, et al., Inadvertent use of bevacizumab in pregnant women with diabetes mellitus type 1, *J Basic Clin Physiol Pharmacol*, 2015;26:161BP3.
53. Sullivan L, Kelly SP, Glenn A, et al., Intravitreal bevacizumab injection in unrecognised early pregnancy, *Eye*, 2014;28:492BP4.
54. Wu Z, Huang J, Sadda S, Inadvertent use of bevacizumab to treat choroidal neovascularisation during pregnancy: a case report, *Ann Acad Med Singapore*, 2010;39:143BP5.
55. Elman MJ, Ayala A, Bressler NM, et al., Intravitreal Ranibizumab for Diabetic Macular Edema with Prompt versus Deferred Laser Treatment: 5BPYear Randomized Trial Results, *Ophthalmology*, 2015;122:375BP81.
56. The economic impact of diabetic macular oedema in Australia. 2015.
57. Lee LJ, Yu AP, Cahill KE, et al., Direct and indirect costs among employees with diabetic retinopathy in the United States, *Curr Med Res Opin*, 2008;24:1549BP59.
58. Mojon B, Pazzi SM, Sousa-Poza A, Mojon DS, Impact of low vision on employment, *Ophthalmologica*, 2010;224:381BP8.
59. Fenwick E, Rees G, Pesudovs K, et al., Social and emotional impact of diabetic retinopathy: a review, *Clin Exp Ophthalmol*, 2012;40:27BP38.
60. Bressler SB, Glassman AR, Almkhatar T, et al., Five-year outcomes of ranibizumab with prompt or deferred laser versus laser or triamcinolone plus deferred ranibizumab for diabetic macular edema, *Am J Ophthalmol*, 2016;164:57–68.